



Effect of plant growth regulators and micronutrients on reproductive attributes of acid lime (*Citrus aurantifolia* Swingle) in *hasta bahar* cropping season

H.K. Deshmukh*, D.H. Paithankar, P.K. Nimbolkar¹, R.K. Dewangan
and C. Awachare¹

Department of Horticulture, P.G.I.
Dr. Panjabrao Deshmukh Krishi Vidyapeeth
Akola - 444104, Maharashtra, India
*E-mail: hdeshmukh975@gmail.com

ABSTRACT

Plant growth regulators and micronutrients at various combinations [GA_3 50ppm + Cycocel 1000ppm + KNO_3 0.2% + Zn 0.3% + Boron 0.1%; GA_3 50ppm + Cycocel 2000ppm + KNO_3 0.2% + Zn 0.3% + Boron 0.1%; GA_3 50ppm + Paclobutrazol 2.5g a.i./tree (soil application) + KNO_3 0.2% + Zn 0.3% + Boron 0.1%; GA_3 50ppm + Paclobutrazol 3.5g a.i./tree (soil application) + KNO_3 0.2% + Zn 0.3% + Boron 0.1%; GA_3 50ppm + Paclobutrazol 1000ppm (foliar application) + KNO_3 0.2% + Zn 0.3% + Boron 0.1%; and, GA_3 50ppm + Paclobutrazol 2000ppm + KNO_3 0.2% + Zn 0.3% + Boron 0.1%] were sprayed before flower emergence in acid lime. Minimum days taken to emergence of flower bud (39.57), duration of flowering (24.07), days to 50% fruit set (6.54) and days taken to fruit maturity (145.90) were observed with application of GA_3 50ppm + Paclobutrazol 3.5g a.i./tree (soil application) + KNO_3 0.2% + Zn 0.3% + Boron 0.1% (T_3), whereas, fruit drop (5.92%) was minimum with GA_3 50ppm + Cycocel 2000ppm + KNO_3 0.2% + Zn 0.3% + Boron 0.1% (T_3). Treatment T_3 also increased the number of flowers per meter length of shoot (49.65) as well as fruit yield (8.90).

Key words: Acid lime, growth regulators, nutrients, flower set, fruit set

INTRODUCTION

Acid lime (*Citrus aurantifolia* Swingle) belongs to the family Rutaceae. It originated in India and has a chromosome number of $2n=18$. Among the various types of citrus fruits grown, acid lime occupies about 3.7 per cent of the total area under citrus in the country. The area under acid lime cultivation in Maharashtra alone is 49.30 hectares, with a production of 739.53MT and productivity of 15.0MT (Anon., 2013).

Acid lime flowers thrice a year, i.e., in the months of January-February, June-July and September-October, under Vidharbha condition and these seasons are generally known as *Ambia*, *Mrig* and *Hasta* bahar, respectively. A bulk of the flowering occurs in *Ambia* bahar (60%), followed by *Mrig* bahar (30%) and *Hasta* bahar (10%). Hence, the market is glutted with *Ambia* bahar fruits, that are harvested in June-July, leading to the lowest price of the fruit during a year in this period. *Hasta* bahar flowering occurs in October-November, and the fruits are ready for harvest in March-May, which is predominately the off-season for acid lime fruits (Thirugananavel *et al*, 2007).

In Vidharbha region, it is highly difficult to regulate *bahar* treatment during September-October due to the absence of sufficient rains. Here, manipulation of *Hasta* bahar flowering with the use of plant growth regulators and other chemicals can serve as an alternative to realize maximum yields during summer, which fetch 6 to 8 times the price of *Ambia* bahar, and 3-4 times the price of *Mrig* bahar season. Keeping the above in view, the present study was aimed at investigating the effect of different combinations of plant growth regulators and micronutrients on reproductive attributes in acid lime in the *Hasta* bahar season.

MATERIAL AND METHODS

A field experiment was conducted in Randomized Block Design (RBD), with three replications, during the year 2013-2014 at the acid lime orchard, College of Horticulture, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola, Maharashtra, situated at 307-457m above mean sea level, of 20.42°N latitude and 72.02°E longitude. Uniform-sized trees were selected, and the required dose of manures, fertilizers, irrigation and other plant protection measures

¹Division of Fruit Crops, ICAR-Indian Institute of Horticultural Research, Hesaraghatta Lake Post, Bengaluru – 560089, Karnataka, India

were applied. Treatment combinations used included: T₁ (Control), T₂ (GA₃ 50ppm + Cycocel 1000ppm + KNO₃ 0.2% + Zn 0.3% + Boron 0.1%), T₃ (GA₃ 50ppm + Cycocel 2000ppm + KNO₃ 0.2% + Zn 0.3% + Boron 0.1%), T₄ [GA₃ 50ppm + Paclobutrazol 2.5g a.i./tree (soil application) + KNO₃ 0.2% + Zn 0.3% + Boron 0.1%], T₅ [GA₃ 50ppm + Paclobutrazol 3.5g a.i./tree (soil application) + KNO₃ 0.2% + Zn 0.3% + Boron 0.1%], T₆ [GA₃ 50ppm + Paclobutrazol 1000ppm (foliar application) + KNO₃ 0.2% + Zn 0.3% + Boron 0.1%] and T₇ [GA₃ 50ppm + Paclobutrazol 2000ppm + KNO₃ 0.2% + Zn 0.3% + Boron 0.1% (foliar application)].

In all the above treatments, water stress for one month was improved from 15th September to 15th October 2013. Spray of plant growth regulators and micronutrients was scheduled as: (i) GA₃ was applied in the first fortnight of June, (ii) Cycocel and Paclobutrazol were sprayed at release of trees from water stress (i.e., 15th September), (iii) Spray of KNO₃, zinc and boron 2 to 3 days prior to release of water stress (i.e., 15th October). Observations were recorded on days taken to emergence of flower-bud, number of flowers per meter length of shoot, duration of flowering, days required for 50% fruit set, fruit drop (%), days to fruit maturity, and yield (t/ha).

RESULTS AND DISCUSSION

Days required for emergence of flower-bud in acid lime was significantly influenced by plant growth regulators and micronutrients (Table 1). After imposition of the treatments, minimum number of days required for emergence of flower-bud (39.57) was observed in treatment T₅ [GA₃ 50ppm + Paclobutrazol 3.5g a.i./tree (soil application) + KNO₃ 0.2% + Zn 0.3% + Boron 0.1%], which

was significantly superior to all the other treatments. Maximum number of days taken for emergence of flower-bud (54.67) was seen in treatment T₁. Similar results were obtained earlier in acid lime, treated with micronutrients and Paclobutrazol (Baskaran *et al*, 2010; Haribabu and Rajput 1982). Similarly, Khanduja *et al* (1974) observed in grape-vine, that zinc promoted nucleic acid synthesis, in turn influencing the formation of flower-bud primordia. Arora (1969) also observed earlier flowering in guava with application of zinc. It is clear from Table 1 that the number of flowers per meter length of shoot was significantly influenced by growth regulators and micronutrients in our study on acid lime. Maximum (49.65) number of flowers per meter length of shoot was observed in treatment T₃ which was significantly superior to all the other treatments, followed by T₅ (44.25) and T₂ (42.63), which were at par with each other. The lowest number of flowers per meter length of shoot (21.40) was recorded in Control (T₁).

Duration of flowering in acid lime under various treatments in our study was found to be non-significant (Table 1). Kachru *et al* (1971) demonstrated that gibberellic acid inhibited flowering in mango, as, higher levels of GA₃ are antagonistic to formation of flower primordia; high levels of endogenous auxins and low levels of endogenous gibberellin favour flower-bud initiation. Paclobutrazol and Cycocel are known inhibitors of gibberellin biosynthesis. Reduced levels of endogenous gibberellins resulting from action of these chemicals may favour early and profuse flowering in mango (Kurian and Iyer, 1993). Growth retardants (Paclobutrazol and Cycocel) have anti-gibberellin effect which, in turn, checks vegetative growth and promotes early flowering. Results depicted in Table 1 reveal that the

Table 1. Effect of different concentrations/ combinations of GA₃, Cycocel, KNO₃, zinc, boron and Paclobutrazol on flowering parameters, fruit set, fruit drop and fruit maturity of acid lime (*Citrus aurantifolia* swingle).

Treatment	Days taken to emergence of flower bud	Flowers per meter length of shoot	Duration of flowering	Days to 50% fruit set	Fruit drop %	Days taken to fruit maturity	Yield kg/plant
T ₁	54.67	21.40	31.33	11.05	12.36 (3.52)	164.50	11.64
T ₂	46.87	42.63	29.33	9.36	6.85 (2.62)	156.70	30.58
T ₃	44.56	49.65	27.67	8.40	5.92 (2.43)	154.40	32.29
T ₄	42.77	40.20	26.45	7.68	8.62 (2.94)	149.60	25.63
T ₅	39.57	44.25	24.07	6.54	7.96 (2.82)	145.90	31.12
T ₆	51.08	32.40	30.87	10.88	9.25 (3.04)	161.30	20.43
T ₇	49.12	36.40	30.45	9.48	9.03 (3.00)	157.70	24.41
'F'-test	*	*	NS	*	*	*	*
SE(m)±	1.44	1.00	1.56	0.51	0.25	2.88	0.66
CD (<i>P</i> =0.05)	4.46	3.08	4.82	1.59	0.80	8.89	2.04

NS = Non-significant; *

number of days required for 50% fruit set in acid lime under various treatments was significantly influenced by plant growth regulators and micronutrients. Treatment T₅ recorded minimum (6.54) number of days for 50% fruit-set, followed by T₄ (7.68) and T₃ (8.40), while, maximum number of days taken for 50% fruit-set was recorded in treatment T₁ (11.05). Application of Cycocel and Paclobutrazol (known anti-gibberellins) cause effective translocation of carbohydrates besides causing a positive effect of cytokinins and auxins in conversion of vegetative buds to flower buds. Similar results were reported in lemon (Monselise *et al*, 1966), mango (Maiti *et al*, 1971) and 'Eureka' lemon (Nir *et al*, 1972). Greenburg *et al* (1993) observed higher fruiting in orange cv. 'Shamouti' on application of Paclobutrazol.

Data depicted in Table 1 reveal that fruit drop in acid lime is significantly influenced by plant growth regulators and micronutrients. Minimum fruit drop was recorded in treatment T₃ (5.92%), which was significantly superior to all the other treatments, followed by T₂ (6.85%), T₅ (7.96%) and T₇ (9.03%). Treatment T₁ (12.36%) showed maximum fruit drop. Haribabu and Rajput (1979) reported in *kagzi* lime that spraying zinc sulphate inhibited abscission, thereby reducing flower and fruit drop. The number of days required to fruit maturity in acid lime was found to be significantly influenced by plant growth regulators and micronutrients (Table 1). Significantly lower number of days (145.90) were required for fruit maturity in treatment T₅, which was significantly superior to all the other treatments. Treatments T₃ (154.40 days), T₂ (156.70 days) and T₇ (157.70 days) were at par with each other. However, treatment T₁ (Control) took maximum number of days for fruit maturity (164.50).

Table 2. Details of treatments applied

Treatment	Treatment details
T ₁	Control
T ₂	GA ₃ 50ppm + Cycocel 1000ppm + KNO ₃ (0.2%) + Zn (0.3%) + Boron (0.1%)
T ₃	GA ₃ 50ppm + Cycocel 2000ppm + KNO ₃ (0.2%) + Zn (0.3%) + Boron (0.1%)
T ₄	GA ₃ 50ppm + Paclobutrazol 2.5g a.i./tree (soil application) + KNO ₃ (0.2%) + Zn (0.3%) + Boron (0.1%)
T ₅	GA ₃ 50ppm + Paclobutrazol 3.5g a.i./tree (soil application) + KNO ₃ (0.2%) + Zn (0.3%) + Boron (0.1%)
T ₆	GA ₃ 50ppm + KNO ₃ (0.2%) + Zn (0.3%) + Boron (0.1%) + Paclobutrazol 1000ppm (foliar application)
T ₇	GA ₃ 50ppm + KNO ₃ (0.2%) + Zn (0.3%) + Boron (0.1%) + Paclobutrazol 2000ppm (foliar application)

Paclobutrazol advances flowering, and ultimately results in early maturity of the fruits.

Significantly high yield (8.90t/ha) was obtained in treatment T₃, followed by T₅ (8.62 t/ha), which was significantly superior to all the other treatments; minimum yield (3.22 t/ha) was seen in T₁ (Control). Appropriate combinations and concentrations of growth regulators and micronutrients (GA₃ 50ppm+ Cycocel 2000ppm + KNO₃ 0.2% + Zinc 0.3% + Boron 0.1%, and utilization of these at the appropriate stage, along with water stress, is effective in regulation of *Hasta* bahar. This is achieved by maximizing the number of fruits and increasing fruit yield harvested during March-April when the demand is higher and prices are at an all-time high. Based on the above findings, appropriate combinations and concentrations of growth regulators, and, micronutrients, i.e., GA₃ 50ppm+Cycocel 2000ppm + KNO₃ 0.2% +Zinc 0.3% + Boron 0.1%, along with imposition of water stress, was found by us to be effective.

This can be very well achieved by applying T₃ and T₅ for increasing fruit yield to be harvested during March - April. Thus, timely application of the treatments GA₃ 50ppm + Cycocel 2000ppm + KNO₃ 0.2% + Zn 0.3% + Boron 0.1% (T₃), and, GA₃ 50ppm + Paclobutrazol 3.5g a.i./tree (soil application) + KNO₃ 0.2% + Zn 0.3% + Boron 0.1% (T₅) promise higher returns to the acid lime grower.

REFERENCES

- Anonymous. 2013. Area and Production of Citrus in India and Maharashtra. <http://www.nhm.gov.in>
- Arora, J.S. 1969. Nutritional studies in mango and guava. Ph.D. thesis submitted to Banaras Hindu University, Varanasi, Uttar Pradesh, India
- Baskaran, A., Renuga, R. and Saraswathy, S. 2011. Advancement of flowering in acid lime by soil applied paclobutrazol. *Madras Agri. J.*, **97**:388-389
- Greenburg, J.E., Goldschidt, E. and Goren, R. 1993. Potential and limitation of the use of PBZ in citrus orchard in Israel. *Acta Hort.*, **329**:58-61
- Haribabu, R.S. and Rajput, C.B.S. 1982. Effect of zinc, 2, 4-D and GA₃ on flowering in *kagzi* lime. *Punjab Hort. J.*, **22**(3-4):140-144
- Kachru, R.B., Susan, R.N. and Guam, E.K. 1971. Inhibition of flowering in mango (*Mangifera indica* L.) by gibberellic acid. *Hort. Sci.*, **6**:140-141
- Khanduja, S.D., Balasubramanyan, V.R. and Saraswat, J.B.

1974. Zinc improves fruit set in grapes. *Indian Hort.*, **19**(4):36
- Kurian, R.M. and Iyer, C.P.A. 1993. Chemical regulation of tree size in mango (*Mangifera indica* L.) cv. Alphonso. I. Effects of growth retardant treatments on vegetative growth and tree vigour. *J. Hortl. Sci. & Biotechnol.*, **68**:349-354
- Maiti, S.C., Mukhopadhyay, A.K. and Sen, P.K. 1971. Effect of growth retardants on flowering and apical dominance of mango. *Curr. Sci.*, **40**(14):388
- Monselise, S.P., Goren, R. and Halevy, A.H. 1966. Effect of B-nine, Cycocel and BTOA on flower bud induction of lemon trees. *Proc. Amer. Soc. Hortl. Sci.*, **89**:195-200
- Nir, I., Goren, R. and Leshem, B. 1972. Effect of water, GA₃ and CCC on flower differentiation in 'Eureka lemon' trees. *J. Amer. Soc. Hortl. Sci.*, **97**(6):774-778
- Thirugananavel, A.R., Amula, W., Baby Rani, K., Indira, P., Mareeswari and S. Parthiban. 2007. Studies on regulation of flowering in acid lime. *Res. J. Agril. Biol. Sci.*, **3**(4):239-241

(MS Received 29 September 2015, Revised 06 June 2016, Accepted 08 June 2016)