



Short communication

## Standardization of recipes for tamarind paste and squash

Archana Pattar, Laxman Kukanoor and Praveen Jholgiker<sup>1</sup>

Department of Post Harvest Technology  
Kittur Rani Channamma College of Horticulture  
Arabhavi - 591 310, India  
E-mail: archana271989@gmail.com

### ABSTRACT

An attempt was made to use tamarind pulp for preparation of paste and squash. Changes in the chemical constituents during storage at ambient temperature were studied. Results showed that in the stored products, TSS, titratable acidity and sugar content increased, whereas, the amount of ascorbic acid decreased. Better quality of paste was prepared from 100g paste + 20g salt with 10ml oil + 20mg BHA, followed by 100g paste + 15g salt with 10ml oil + 20mg BHA. Squash with 30% juice, 50% TSS, 1% acidity and 0.5% salt was superior to the other recipes. Both tamarind paste and squash retained their characteristic colour, aroma, taste and were microbiologically safe up to 3 months of storage at room temperature.

**Key words:** Tamarind, pulp, paste, salt, spiced squash, storage, and nutritional quality

Tamarind (*Tamarindus indica* L.) is an arboreal fruit of the family Fabaceae. The name 'tamarind' is derived from the Arabic term 'tamar-ul-Hind' which means 'date of India'. The most outstanding characteristic of tamarind is its acidity, with total acidity ranging from 12.2% to 23.8% as tartaric acid. Tamarind is a good source of Vitamin B (Siddig *et al*, 2006). The fruit is used for making different products such as powder, juice, concentrate, paste, kernel powder, tartaric acid, pectin, tartarates and alcohol. Tamarind products have been extensively used in traditional Indian and African medicine. Although several medicinal values are claimed for various preparations from the fruit, leaf, flower, bark, etc. of the tamarind tree, only antiscorbutic properties of the pulp, laxative action of the juice and diuretic properties of the leaf sap are well-established (Siddig *et al*, 2006).

Looking into the fast increasing area under tamarind cultivation, methods of preservation of tamarind products need to be developed to regulate prices of the fresh tamarind fruit and to safeguard interest of the tamarind grower during a period of glut (Joshi *et al*, 2012). Nowadays, people prefer 'instant' preparations and, hence, tamarind could be more easily used in the form of paste. Value-added products such as tamarind paste, with proper packaging, can be easily transported to areas where it is difficult to transport the pulp. Tamarind products possess good export potential (Champakam and Peter, 2000). Due to the perishable nature

of fruits, these require immediate processing, to avoid post-harvest losses (Ramakumar *et al*, 1997) and changes in colour from brown to black due to phenolics, and non-enzymatic browning during storage. Therefore, the present investigation was carried out to develop a technology for preparation of value-added products (paste and squash) in tamarind.

Fully ripe tamarind fruits of cv. DTS-1 were procured from Department of Plantation and Spices Crops, Kittur Rani Channamma College of Horticulture, Arabhavi. The pulp was manually separated from the shell and seed. Ginger (*Zingiber officinalis*) and mango ginger (*Curcuma amada*) were procured from the market.

The experiment, with seven treatments, was conducted using Completely Randomized Block design. Tamarind paste was prepared by adding different concentration of salt, and the squash was prepared by varying tamarind juice concentration and the TSS.

### Treatment details

#### Recipe for tamarind paste

T <sub>1</sub> - 100g paste + 5g salt	T <sub>5</sub> - 100g paste + 25g salt
T <sub>2</sub> - 100g paste + 10g salt	T <sub>6</sub> - 100g paste + 30g salt
T <sub>3</sub> -100g paste + 15g salt	T <sub>7</sub> - 100g paste (Control) without salt
T <sub>4</sub> -100g paste + 20g salt	

<sup>1</sup>Dept. of Fruit Science, KRC College of Horticulture, Arabhavi, Karnataka, India

Sunflower oil (10ml) and BHA (Butylated hydroxy anisole) were added (20mg) in all the treatments

**Recipe for tamarind squash blended with ginger: all the treatments had identical acidity (1%), salt (0.5%) and KMS (0.25g/l)**

Treatment	Juice (%)	TSS (%)
T <sub>1</sub>	25 (9:1) Tamarind: Ginger	40
T <sub>2</sub>	30 (9:1) Tamarind: Ginger	40
T <sub>3</sub>	25 (9:1) Tamarind: Ginger	45
T <sub>4</sub>	30 (9:1) Tamarind: Ginger	45
T <sub>5</sub>	25 (9:1) Tamarind: Ginger	50
T <sub>6</sub>	30 (9:1) Tamarind: Ginger	50
T <sub>7</sub>	30 (9:1) Tamarind : Mango ginger	40

**Preparation of tamarind paste:** One kg of de-seeded pods were taken and soaked in water in the ratio 1:2 for four hours. The juice was extracted by repeated squeezing of the soaked pulp and by passing it through one mm sieve. Juice so extracted was boiled until it attained the required consistency. The end-point was judged by recording the TSS (32-34° Brix). At the end of boiling, table salt, edible oil and BHA were added as per treatments shown above, and mixed thoroughly. Sunflower oil is a preservative and BHA is an antioxidant. Then, the product was filled while hot into a pre-sterilized PET jar to hold 200g, and sealed. The jars were stored at room temperature (28-38°C) three months.

**Preparation of tamarind squash:** Tamarind juice was mixed with sugar and water as per recipes mentioned in treatment details. The squash was preserved by adding potassium metabisulphate @ 0.25 gram per litre and salt at 0.5%. Prepared squash was filled in clean, sterile bottles of 200ml capacity and sealed. This was stored at room temperature (28-38°C) for three months.

**Chemical analysis:** TSS was measured using a digital refractometer (Make:Erma, Japan). pH meter was used to measure the pH (Model: Analog Research, USA). Titratable acidity was estimated as per the modified procedure of AOAC (Anon., 1984). Ascorbic acid content was determined titrimetrically using 2, 6-dichlorophenol indophenol dye as per the modified procedure of AOAC (Anon., 1984). Sugars were estimated by dinitro salicylic acid (DNSA) method (Miller, 1972).

Organoleptic evaluation of tamarind squash was done at 30 day intervals, i.e., 30, 60 and 90 days after storage

(DAS). The squash was diluted with water in the ratio 1:3 before serving. Organoleptic characters like colour and appearance, aroma and flavour, taste and overall acceptability were evaluated by a panel of semi-trained judges, on a five-point Hedonic scale (Ranganna, 1997).

**Microbial analysis:** Microbial count of the tamarind beverages after storage was taken as per Harrigan and Mccance (1996).

**Preparation of sample:** Samples were prepared by taking 10ml of the representative sample from each treatment. Each sample was mixed with 90ml sterilized distilled water in a conical flask.

**Dilution:** Serial dilution technique was applied to estimate bacterial, fungal and yeast load in the tamarind products. Dilutions 10<sup>-5</sup> and 10<sup>-6</sup> were used for counting bacteria and 10<sup>-3</sup> and 10<sup>-4</sup> were used for counting fungi and yeast. Nutrient Agar media, Rose Bengal agar media and Wichraham media were used to enumerate bacterial, fungal and yeast colonies, respectively, in the processed products of tamarind.

**Enumeration:** The medium was sterilized in a pressure cooker at 121°C for 20 minutes. Into each sterilized petri dish, 1ml of the respective aliquot was transferred; 10ml of the medium was poured into duplicate plates. After solidification, the plates were incubated upside down at 37°C for six days.

**Colony count:** The colonies were counted and the total count was expressed by multiplying the number of colonies with the dilution factor, and expressed as CFU per gram or ml of the sample.

**Changes in chemical composition of tamarind paste during storage:** Significantly high TSS was found in T<sub>6</sub> consisting of 100g paste + 30g salt + 10ml oil + 20mg BHA at 30, 60 and 90 days after storage. These differences between treatments may be attributed to different levels of the salt used. Difference in the TSS has been seen to be mainly due to the addition of different concentration of salt by Kotecha and Kadam (2003). Highest pH was recorded in T<sub>7</sub> at 30, 60 and 90 days after storage. Corresponding decrease in acidity might be responsible for high pH. Similar observations for change in pH were made by Nath *et al* (2005) in kinnow mandarin-ginger squash (Table 1). Significantly high titratable acidity was recorded in the Control (Fig. 1). Decrease in acidity during storage could be attributed to chemical interaction induced by temperature between chemical constituents of the juice and the action of enzymes (Mapson, 1970).

With increase in storage period, ascorbic acid content decreased (Fig 2). Significantly high ascorbic acid content remained in T<sub>6</sub> consisting of 100g paste + 30g salt + 10ml oil + 20mg BHA. Gradual decrease seen in ascorbic acid content may be due to oxidative destruction of this acid by ascorbic acid enzymes in the presence of molecular oxygen and, may also be due to the effect of storage temperature and catalytic activity of fructose during storage (Mapson, 1970). Higher amount of reducing and non-reducing sugars were recorded in T<sub>6</sub> and T<sub>5</sub> at 30, 60 and 90 DAS. At 30 days after storage significantly high total sugars were recorded in T<sub>6</sub> (29.55 %). However, total sugars at 60 or 90 days after storage were found non-significant. Increase seen in sugar content could be due to the slow hydrolysis of polysaccharides, acids and pectic substances into simpler substances like sugars (Table 2). These findings are in accordance with Sarita *et al* (2006) in Rangpur lime RTS and squash, and Kichu (2008) in sapota beverages.

**Table 1. Changes in TSS and pH of tamarind paste during storage**

Treatment	TSS (%)*			pH**		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T <sub>1</sub> - 100g paste + 5g salt	34.10	29.56	30.40	2.87	2.96	3.12
T <sub>2</sub> - 100g paste + 10g salt	32.83	33.60	35.23	2.85	2.94	3.01
T <sub>3</sub> - 100g paste + 15g salt	33.60	35.82	37.16	2.74	2.83	2.97
T <sub>4</sub> - 100g paste + 20g salt	35.10	39.82	40.23	2.48	2.57	2.73
T <sub>5</sub> - 100g paste + 25g salt	34.80	35.71	37.86	2.71	2.80	2.95
T <sub>6</sub> - 100g paste + 30g salt	38.50	41.85	42.60	2.51	2.60	2.77
T <sub>7</sub> - 100g paste (control)	32.10	25.08	23.03	3.10	3.19	3.35
CD ( <i>P</i> =0.01)	0.553	4.433	1.917	0.041	0.024	0.041

\*Initial TSS content was 34% DAS - Days after storage

\*\* Initial pH was 3.0

**Table 2. Changes in various sugars in tamarind paste during storage**

Treatment	Reducing sugars (%)*			Non-reducing sugars (%)**			Total sugars (%)***		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T <sub>1</sub> - 100g paste + 5g salt	22.46	22.77	23.09	2.88	3.07	3.25	25.34	25.84	26.34
T <sub>2</sub> - 100g paste + 10g salt	23.63	23.94	24.24	2.10	2.29	2.49	25.73	26.23	26.73
T <sub>3</sub> - 100g paste +15g salt	24.23	24.54	24.84	2.47	2.66	2.86	26.70	27.20	27.70
T <sub>4</sub> - 100g paste +20g salt	25.03	25.35	25.65	3.07	3.25	3.45	28.10	28.60	29.10
T <sub>5</sub> - 100g paste +25g salt	22.63	22.95	23.25	3.77	3.95	4.15	26.40	26.90	27.40
T <sub>6</sub> - 100g paste +30g salt	26.93	27.25	27.55	2.62	2.80	3.00	29.55	30.05	30.55
T <sub>7</sub> - 100g paste (Control)	21.05	21.36	21.67	3.18	3.47	3.66	24.23	24.83	25.33
CD ( <i>P</i> =0.01)	0.037	0.048	0.024	1.027	0.240	0.249	1.057	NS	NS


\*Initial reducing sugars 20.02% DAS - Days after storage

\*\*Initial non-reducing sugars 2.20% NS- Non Significant

\*\*\*Initial total sugars 22.22%

**Changes in chemical composition of tamarind squash during storage:**


An increasing trend was observed in the content of total soluble solids in tamarind-ginger blended squash during storage period (Table 3). Treatment T<sub>6</sub> recorded highest TSS compared to other treatments during storage which could be due to hydrolysis of polysaccharides



**Fig 1. Changes in titratable acidity (%) of tamarind paste during storage.**

- T<sub>1</sub>- 100g paste + 5g salt
- T<sub>2</sub>- 100g paste + 10g salt
- T<sub>3</sub>-100g paste + 15g salt
- T<sub>4</sub>-100g paste + 20g salt
- T<sub>5</sub>- 100g paste + 25g salt
- T<sub>6</sub>- 100g paste + 30g salt
- T<sub>7</sub>- 100g paste (Control)

\*Initial titratable acidity was 12.5%



**Fig 2. Changes in ascorbic acid content (mg/100g) of tamarind paste during storage.**

- T<sub>1</sub>- 100g paste + 5g salt
- T<sub>2</sub>- 100g paste + 10g salt
- T<sub>3</sub>-100g paste + 15g salt
- T<sub>4</sub>-100g paste + 20g salt
- T<sub>5</sub>- 100g paste + 25g salt
- T<sub>6</sub>- 100g paste + 30g salt
- T<sub>7</sub>- 100g paste \*(Control)

\*Initial ascorbic acid content was 4.00mg/100g

like starch and pectic substances into simpler substances. Similar observations were recorded by Kotecha and Kadam (2003) in tamarind syrup and Nath *et al* (2005) in ginger-blended kinnow mandarin squash. Among the treatments, T<sub>1</sub> recorded the highest pH, which was on par with T<sub>2</sub> (2.70) and T<sub>7</sub> (2.62, 2.64 and 2.75 corresponding to 30, 60 and 90

**Table 3. Changes during storage in TSS and pH of tamarind squash blended with ginger**

#Treatment	TSS (%)*			pH **		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T <sub>1</sub> - 25% juice + 40% TSS	40.16	40.33	40.96	2.74	2.75	2.79
T <sub>2</sub> - 30% juice + 40% TSS	40.33	40.70	41.70	2.70	2.70	2.76
T <sub>3</sub> - 25% juice + 45% TSS	45.16	45.76	46.40	2.32	2.45	2.47
T <sub>4</sub> - 30% juice + 45% TSS	47.10	47.53	48.20	2.24	2.26	2.32
T <sub>5</sub> - 25% juice + 50% TSS	53.23	53.40	54.40	2.16	2.21	2.41
T <sub>6</sub> - 30% juice + 50% TSS	53.70	54.03	54.66	2.13	2.16	2.16
T <sub>7</sub> - 30% juice + 40% TSS	41.20	41.33	41.70	2.62	2.64	2.75
CD ( <i>P</i> =0.01)	0.265	0.175	1.178	0.263	0.280	0.360

# Acidity (1%) and salt (0.5%) \* Initial TSS in T<sub>1</sub> T<sub>2</sub> and T<sub>7</sub> = 40%  
DAS - Days after storage in T<sub>3</sub> and T<sub>4</sub> = 45%  
in T<sub>5</sub> and T<sub>6</sub> = 50%

\*\* Initial pH = 2.0

**Table 4. Changes during storage in titratable acidity and ascorbic acid content of tamarind squash blended with ginger**

#Treatment	Titratable acidity (%)*			Ascorbic acid (mg/100g)**		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T <sub>1</sub> - 25% juice + 40% TSS	0.97	0.95	0.93	3.73	3.64	3.60
T <sub>2</sub> - 30% juice + 40% TSS	0.96	0.94	0.92	3.75	3.70	3.62
T <sub>3</sub> - 25% juice + 45% TSS	0.93	0.92	0.89	3.71	3.61	3.56
T <sub>4</sub> - 30% juice + 45% TSS	0.92	0.91	0.88	3.58	3.56	3.54
T <sub>5</sub> - 25% juice + 50% TSS	0.91	0.89	0.86	3.40	3.28	3.20
T <sub>6</sub> - 30% juice + 50% TSS	0.90	0.87	0.84	3.67	3.64	3.51
T <sub>7</sub> - 30% juice + 40% TSS	0.95	0.94	0.91	3.71	3.62	3.58
CD ( <i>P</i> =0.01)	0.024	0.022	0.028	0.102	0.067	0.264

# Acidity (1%) and salt (0.5%)

DAS - Days after storage

\*Initial titratable acidity = 1.0%

\*\*Initial ascorbic acid content = 3.75mg/100

days, respectively). This was mainly due to a corresponding decrease in acidity. Similar observation for change in pH was reported by Nath *et al* (2005) in kinnow mandarin-ginger squash (Table 3).

Acidity of the products prepared gradually declined during storage (Table 4). Among the various treatments, maximum acidity was noticed in T<sub>1</sub> which was on par with T<sub>2</sub>. This decrease in acidity may have been due to acid hydrolysis of polysaccharides and non-reducing sugars into their simpler components, where, the acid is utilized for converting the former into hexose sugars or complexes in the presence of metal ions. Analogous result was reported by Gajanana (2002) in amla juice. Reduction in acidity during storage period in beverages was observed by Lakshmi *et al* (2005) in flavoured tamarind RTS beverages and Nidhi *et al* (2008) in RTS bael-guava beverage. Gradual decrease during storage in ascorbic acid content in tamarind squash blended with ginger was observed (Table 4). Among the treatments, T<sub>2</sub> (3.68mg/100g), which was on par with T<sub>1</sub> (3.64mg/100g) and T<sub>6</sub> (3.60mg/100g), recorded maximum ascorbic acid retention during storage compared to other treatments. Decrease in ascorbic acid content could be due to the effect of storage temperature (Mapson, 1970). As ascorbic acid content of a squash is directly dependent on blended-juice used for its preparation, comparatively higher ascorbic acid content was observed at the end of storage period in the case of samples with higher percentage of juice, in all the treatments. Analogous observations for decline in ascorbic acid content were reported in aonla juice by Gajanana (2002), in rose apple-aonla squash by Basavaraja (2005), and in bael-guava RTS by Nidhi *et al* (2008).

Among the treatments, the maximum amount of reducing sugars, non-reducing sugars and total sugars were noticed in T<sub>6</sub> consisting of 30% juice + 50% TSS (Table 5). Total sugar content of tamarind-blended ginger squash increased only slightly during storage. This could be attributed to acid hydrolysis of polysaccharides that result in an increase in soluble sugars. Increase in reducing and total sugars, and decrease in non-reducing sugars, is a general phenomenon as has been noticed by several workers. Kotecha and Kadam (2003) in tamarind syrup and Sahu *et al* (2006) in mango-lemongrass beverage observed an increase in total and a reducing sugars and decrease in non-reducing sugars during storage.

**Organoleptic evaluation of tamarind squash (scores out of 5.00):** For 30, 60 and 90 days after storage of

**Table 5. Changes during storage in sugars in tamarind squash blended with ginger**

Treatment#	Reducing sugars (%)			Non-reducing sugars (%)			Total sugars (%)		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T <sub>1</sub> - 25% juice + 40% TSS	8.53	8.82	9.12	13.74	13.95	14.15	22.27	22.77	23.27
T <sub>2</sub> - 30% juice + 40% TSS	8.63	8.96	9.25	17.67	17.84	18.05	26.30	26.80	27.30
T <sub>3</sub> - 25% juice + 45% TSS	8.80	9.10	9.40	19.50	19.70	19.97	28.30	28.80	29.30
T <sub>4</sub> - 30% juice + 45% TSS	8.94	9.23	9.53	18.39	18.60	18.80	27.33	27.83	28.33
T <sub>5</sub> - 25% juice + 50% TSS	9.08	9.37	9.67	14.32	14.53	14.73	23.40	23.90	24.40
T <sub>6</sub> - 30% juice + 50% TSS	9.22	9.51	9.81	20.53	20.74	20.94	29.75	30.25	30.75
T <sub>7</sub> - 30% juice + 40% TSS	8.62	8.93	9.22	16.61	16.80	17.01	25.23	25.73	26.23
CD ( <i>P</i> =0.01)	0.037	0.024	0.312	0.191	NS	NS	NS	NS	NS

#Acidity (1%) and salt (0.5%)

\*Initial reducing sugars = 8.25%

DAS - Days after storage

\*\*Initial non-reducing sugars = 13.40%

NS- Non significant

\*\*\*Initial total sugars = 21.65%


**Table 6. Organoleptic scores (out of 5.00) for colour, aroma and taste of tamarind squash as influenced by various treatments**

Treatment	Colour score (out of 5.00)			Aroma score (out of 5.00)			Taste score (out of 5.00)		
	30 DAS	60 DAS	90 DAS	30DAS	60 DAS	90 DAS	30DAS	60 DAS	90 DAS
T <sub>1</sub> - 25 % juice + 40% TSS	3.30	3.20	3.18	3.00	2.41	2.21	3.25	3.35	3.46
T <sub>2</sub> - 30 % juice + 40% TSS	3.50	3.40	3.38	3.51	2.83	2.68	3.54	3.61	3.75
T <sub>3</sub> - 25 % juice + 45% TSS	3.40	3.30	3.28	3.16	3.08	2.86	3.38	3.41	3.55
T <sub>4</sub> - 30 % juice + 45% TSS	3.80	3.78	3.76	3.78	3.58	3.25	3.85	3.91	4.05
T <sub>5</sub> - 25 % juice + 50% TSS	3.70	3.68	3.66	3.68	2.98	2.51	3.93	4.18	4.48
T <sub>6</sub> - 30 % juice + 50% TSS	3.90	3.80	3.78	3.83	3.75	3.48	3.90	4.13	4.26
T <sub>7</sub> - 30 % juice + 40% TSS	3.60	3.50	3.45	3.26	3.08	2.66	3.18	3.21	3.35
CD ( <i>P</i> =0.01)	0.24	0.184	0.210	0.551	0.879	0.644	0.426	0.487	0.483

DAS - Days after storage

tamarind squash significantly maximum score for colour, aroma & overall acceptability was observed in treatment T<sub>6</sub>, whereas for taste maximum score was found in treatment T<sub>5</sub> (Table T<sub>6</sub> & Fig 3). This might be due to better consistency, acceptable color and sugar acid blend. Similar result has been reported (Kotecha and Kadam, 2003) in tamarind RTS.

**Microbial load:** Safety of a product with respect to its microbial load is as important as its chemical composition. In tamarind paste, bacterial, fungal and yeast populations were fewer in treatment T<sub>4</sub> ( $2.60 \times 10^5$ ,  $0.90 \times 10^3$  and  $2.4 \times 10^3$  CFU/ml, respectively). Reasons for the low microbial load could be on inhibitory effect of salt and acid on these microorganisms; whereas, in squash, bacterial, fungal and yeast populations were found to be the minimum in treatment T<sub>3</sub> ( $4.3 \times 10^5$ ,  $2.41 \times 10^3$  and  $1.61 \times 10^3$  CFU/ml, respectively), followed by treatment T<sub>6</sub> ( $4.41 \times 10^5$ ,  $2.52 \times 10^3$  and  $1.72 \times 10^3$  CFU/ml, respectively) (Tables 7 and 8). A low microbial load could be due to an inhibitory effect of high sugar concentration (55-70°Brix) and inhibitory effect of acid and

**Fig 4. Changes in overall acceptability of stored tamarind squash blended with ginger.**

T<sub>1</sub>- 25% juice + 40% TSS    T<sub>5</sub>- 25% juice + 50% TSS  
 T<sub>2</sub>- 30% juice + 40% TSS    T<sub>6</sub>- 30% juice + 50% TSS  
 T<sub>3</sub>- 25% juice + 45% TSS    T<sub>7</sub>- 30% juice + 40% TSS  
 T<sub>4</sub>- 30% juice + 45% TSS

KMS on microorganisms. Similar results were also reported by Li *et al* (1989) in orange juice, Shobana (2003) in guava beverages and Kotecha and Kadam (2003) in tamarind concentrate.

**Table 7. Effect of various treatments on microbial load in tamarind squash**

Treatment	Microbial load (No. x 10 <sup>3</sup> CFU/ml) three months after storage		
	Bacteria	Fungi	Yeast
T <sub>1</sub> - 25% juice + 40% TSS	5.06	3.2	2.4
T <sub>2</sub> - 30% juice + 40% TSS	5.22	3.3	2.5
T <sub>3</sub> - 25% juice + 45% TSS	4.64	2.78	1.98
T <sub>4</sub> - 30% juice + 45% TSS	4.88	2.99	2.19
T <sub>5</sub> - 25% juice + 50% TSS	4.3	2.41	1.61
T <sub>6</sub> - 30% juice + 50% TSS	4.41	2.52	1.72
T <sub>7</sub> - 30% juice + 40% TSS	5.08	3.19	2.39

**Table 8. Effect of various treatments on microbial load in tamarind paste**

Treatment	Microbial load (No. x 10 <sup>3</sup> CFU/ml) three months after storage		
	Bacteria	Fungi	Yeast
T <sub>1</sub> - 100g paste + 5g salt	20.75	1.6	67.55
T <sub>2</sub> - 100g paste + 10g salt	10.25	2.20	3.05
T <sub>3</sub> - 100g paste + 15g salt	3.25	1.35	2.90
T <sub>4</sub> - 100g paste + 20g salt	2.60	0.90	2.4
T <sub>5</sub> - 100g paste + 25g salt	4.35	1.0	18.7
T <sub>6</sub> - 100g paste + 30g salt	3.05	1.70	4.2
T <sub>7</sub> - 100g paste (control)	37.15	2.40	115.5

CFU- Colony forming unit

**REFERENCES**

- Anonymous. 1984. Official Methods of Analysis. Sidney Williams (Ed.), Association of Official Analytical Virginia, 14<sup>th</sup> Edition, pp. 424-462
- Basavaraja, P.S. 2005. Value added products from rose apple (*Syzigium jambos* Alston). M.Sc. (Hort.) Thesis, Uni Agric. Sci., Bengalore
- Champakam, B. and Peter, K.V. 2000. Processing and chemistry of tamarind (*Tamarindus indica* Linn.) *Spice India*, **13**:9-12
- Gajanana, K. 2002. Processing of aonla (*Emblica officinalis* Gaertn.) fruits. M.Sc. (Hort.) Thesis, Univ. Agri. Sci., Dharwad, India
- Joshi, A.A., Kshirsagar, R.B. and Sawate, A.R. 2012. Studies on standardization of enzyme concentration and process for extraction of tamarind pulp, variety Ajanta. *J. Food Process and Tech.*, **3**:1-3
- Kichu, T. 2008. Processing and value addition to sapota fruits. M.Sc. (Hort.) Thesis, Univ. Agric. Sci., Dharwad
- Kotecha, P.M. and Kadam, S.S. 2003. Preparation of ready to serve beverage, syrup and concentrate from tamarind. *J. Food Sci. Tech.*, **40**:76-79
- Li, Z., Alli, I. and Kermasha, S. 1989. Use of acidification, low temperature and sarbates for storage of juice. *J. Food Sci.*, **54**:674-678
- Mapson, C.W. 1970. Vitamins in fruits. In: Biochemistry of fruits and their products. Vol. 1. Hulme, A.C. (Ed.), Academic Press, New York, pp. 376-387
- Miller, G.L. 1972. Use of dinitro-salicylic acid reagent for determination of reducing sugar. *Anat. Chem.*, **31**:426-428
- Nath, A., Yadav, D.S., Pranabjoyoti Sarma and Dey, B. 2005. Standardization of ginger-kinnow squash and its storage. *J. Food Sci. Tech.*, **42**:520-522
- Nidhi, Rakesh Gehlot, Singh, R. and Rana, M.K. 2008. Changes in chemical components of RTS bael-guava blended beverages during storage. *J. Food Sci. Tech.*, **45**:378-380
- Ramakumar, M.V., Babu, C.K., Subramanya, S., Ranganna, B. and Krishnamurthy, K.C. 1997. Development of a tamarind de-huller and short-term storage of pulp. Proc. Nat. Sym. on *Tamarindus indica* L. Tirupathi (A.P.), 27-28 June, pp 145-150
- Ranganna, 1997. Manual of analysis of fruit and vegetable products. II Edition. Tata McGraw Hill Publishing Company Ltd., New Delhi, India
- Sahu, C., Choudhary, P.L., Patel, S. And Sahu, R., 2006. Physico-chemical and sensory characteristics of whey based mango herbal (lemongrass) beverage. *Ind. Food Pac.*, **60**:127-132
- Sarita, S., Chopra, C.S. and Santosh Kumari, 2006. Studies on physico-chemical characteristics of Rangpur lime (*Citrus limonia* Osbeck.). *Beverage Food Wld.*, **31**:44-45
- Shobana, N. 2003. Standardization of processing technology for banana fruits. M.Sc. (Hort.) thesis, Univ. Agril. Sci., Dharwad, India
- Siddig, K.E., Gunasena. H.P., Prasad B.A., Pushpakumar, D.K., Ramana. K.V., Vijayanand, P. and Williams, J. T. 2006. Tamarind monograph, Southampton Centre for Underutilized Crops, Southampton, U.K., pp. 1-198

(MS Received 31 January 2013, Revised 16 September 2013, Accepted 28 October 2013)