

Original Research Paper

**Development and evaluation of novel gladiolus hybrid selections
IIHRG-7 (IC620379) and IIHRG-11 (IC620380) for
flower quality and *Fusarium* wilt resistant**

**Manjunatha Rao T., Janakiram T., Negi S.S., Aswath C., Dhananjaya M.V.,
Rajiv Kumar and Ramachandran N.**

Division of Flower and Medicinal Crops, ICAR-Indian Institute of Horticultural Research
Hesaraghatta Lake Post, Bengaluru 560 089, Karnataka, India

*E mail: Rajiv.Kumar11@icar.gov.in

ABSTRACT

The present study was carried out to evaluate the performance of two novel gladiolus hybrid selections IIHRG-7 and IIHRG-11 along with commercial checks, for flower quality traits and *Fusarium* wilt resistant in Completely Randomized Block Design, during 2012-13 to 2014-15. Data of three years were pooled and analyzed statistically. The hybrid selections IIHRG-7 and IIHRG-11 had been developed through hybridization by crossing Arka Meera x Picardy and Gold Medal 412 x Arka Poonam, respectively followed by selection. IIHRG-7 has novel flower colour (as per RHS Colour Chart) *i.e.* Red-Purple (65.B) having Red-Purple (62.A) streaks with Red-Purple (67.B) splash and spike with variegated florets, while, IIHRG-11 has novel floret colour as Red (41.C) having Red (41.A) margin. Blotch Red (46.B) with yellow (13.C) border and resistant to *Fusarium* wilt disease. These hybrid selections are suitable for cut-flower and flower arrangement purposes. Further, these hybrid selections will be useful for developing new gladiolus hybrid selections with novel traits and resistant to *Fusarium* wilt disease.

Key words: Gladiolus, evaluation, flowering, corm, vase life

INTRODUCTION

Gladiolus is one of the most important bulbous flowering plant commercially grown for cut flowers, garden display and floral arrangement. It belongs to the family Iridaceae and sub-family Ixioideae. It ranks second in area (20.53'000 ha) and production (132.58'000 tons) among the cut flowers grown in India (Anonymous, 2016). The main emphasis in gladiolus improvement has to be given on development of varieties having attractive novel colour and more number of well spaced large sized florets mainly for cut flower, long spikes and good corm multiplication ability (Swaroop *et al.*, 2018). *Fusarium* wilt is the most devastating disease in gladiolus which is caused by the fungus *Fusarium oxysporum* Schltdl. f.sp. *gladioli* (Massey) W.C. Snyder & H.N. Hansen (Massey, 1926 and Nelson *et al.*, 1981). It is a major bottleneck in gladiolus cultivation causing 60-80%

crop damage and huge economic loss to flower growers (Lakshman *et al.*, 2012 and Kakadeet *et al.*, 2016).

As conventional management practices for *Fusarium* wilt disease include corm treatment with fungicides and soil fumigation are time consuming, labour intensive and increase the cost of cultivation, developing *Fusarium* wilt disease resistance gladiolus genotypes is an economical viable option in managing this disease. Identification of genetic resources for resistance to *Fusarium* wilt is crucial for harnessing resistance from these plants which can be deployed in development of resistant varieties. Therefore, the present study was carried out to evaluate two novel gladiolus hybrid selections IIHRG-7 and IIHRG-11 for their flower quality and *Fusarium* wilt disease resistance.

MATERIAL AND METHODS

Hybridization followed by selection was employed to develop novel gladiolus hybrid selections IIHRG-7 and IIHRG-11 involving crosses Arka Meera x Picardy and Gold Medal 412 x Arka Poonam during 1986 and 1988, respectively. From hybrid seeds, cormels were produced. After the period of dormancy, cormels were planted and corms were harvested. Promising novel hybrid selections *viz.*, IIHRG-7 and IIHRG-11 were selected and multiplied vegetatively. Further, these hybrid selections with commercial checks Pink Friendship and *Psittacinus* hybrid, were evaluated for flower quality traits and resistance to *Fusarium* wilt disease, in replicated trial in Completely Randomized Block Design for three consecutive years i.e. from 2012-13 to 2014-15. The data on various biometrical parameters recorded were subjected to statistical analysis (Panse and Sukhatme, 1967).

Screening for resistance was undertaken in pot culture inside polyhouse in replicated trial using Completely Randomized Block Design. Uniform sized corms (5.5

cm to 6 cm) of IIHRG-11 and Pink Friendship (check) were planted in plastic pots containing 2 kg sterilized growing media @ 2:1:1 v/v (soil: sand: FYM). Sorghum based *Fusarium oxysporum* f. sp. gladioli inoculum was mixed in the soil at 4 g per 100 g of soil one day before planting and watering was done. The response of genotypes to *Fusarium* inoculation was evaluated at 90 days after planting (Elewa *et al.*, 2001). The disease incidence in per cent was recorded according to Riaz *et al.* (2010) and categorization of gladiolus genotypes based on disease incidence percentage was carried out as instructed by Shanmugam *et al.* (2009) as follows: 0-10% = Highly resistant (HR); 10-25% = Resistant (R); 25-50% = Moderately susceptible (MS); 50-75% = Susceptible (S); 75-100% = Highly susceptible (HS). The results have been presented and discussed at the probability level of one per cent. The data regarding disease incidence and mortality were recorded using following formulae:

$$\text{Disease incidence per cent} = \frac{\text{Number of diseased plants}}{\text{Total number of plants}} \times 100$$

$$\text{Mortality per cent} = \frac{\text{Number of plants died due to disease}}{\text{Total number of plants}} \times 100$$

RESULTS AND DISCUSSION

On the perusal of data presented in Table 1 indicated significant differences between hybrid selection IIHRG-7 and Pink Friendship (check) for most of the vegetative and floral characters, however, plant height, floret diameter, number of marketable spikes per corm and vase life were found non-significant. The genotype Pink Friendship (check) recorded significantly early spike emergence (53.00 days) and flowering (62.67 days) in comparison to IIHRG-7 (63.95 days and 72.25 days, respectively). IIHRG-7 recorded significantly higher spike length (123.11 cm) than the check Pink Friendship (113.70 cm), however, longest rachis was recorded in Pink Friendship (57.41 cm). The spike length is one of the major criteria in selection of superior hybrid selection in gladiolus. The Pink Friendship (check) recorded more number of florets per spike (17.10) and florets

remain open at a time (6.12) than IIHRG-7 (12.66 and 5.55, respectively). However, IIHRG-7 recorded significantly maximum number of total spikes per corm (1.66) than the Pink Friendship (1.40). The more number of spikes per corm are directly related to the higher productivity per unit area. The Pink Friendship recorded higher flowering duration (12.03 days) than the IIHRG-7 (9.61 days) owing to presence of more number of florets per spike in Pink Friendship which opened in acropetal successions for longer period. Sankari *et al.* (2012) reported variation in flowering traits in 42 gladiolus genotypes and recommended genotypes Pusa Swarnima, Pusa Shagun, Thumbolina, Priscilla and Candyman for cut flower production under Eastern Ghats of Tamil Nadu. Safeena and Thangam (2019) also evaluated ten cultivars of gladiolus for flowering traits and recommended Arka Amar and Darshan for cut flower purpose under Goa conditions.

Table 1. Vegetative and floral traits of Gladiolus hybrid selection IHRG-7 with check Pink Friendship (pooled data of three years)

Genotype	Days to spike emergence	Days to flower	Plant height (cm)	Spike length (cm)	Rachis length (cm)	Floret diameter (cm)	No. of florets per spike	Florets remain open at a time	No. of spikes/corm	No. of marketable spikes/corm	Flowering duration (days)	Vase life (days)
IHRG-7	63.95	72.25	142.06	123.11	46.77	10.66	12.66	5.55	1.66	1.43	9.61	9.33
Pink Friendship	53.00	62.67	144.88	113.70	57.41	10.79	17.10	6.12	1.40	1.40	12.03	9.00
C.D. at 5%	3.01	9.67	NS	6.23	2.97	NS	0.97	0.41	0.28	NS	1.11	NS

Table 2. Corm and cormels traits of Gladiolus hybrid selection IHRG-7 with check Pink Friendship (pooled data of three years)

Genotype	Corm per corm (Nos.)	Cormel per corm (Nos.)	Diameter of corm (cm)	Diameter of cormel (cm)	Weight of corm (g)	Weight of cormel (g)
IHRG-7	1.46	26.63	6.75	1.17	72.44	0.57
Pink Friendship	1.03	52.11	5.65	1.02	61.33	0.44
C.D. at 5%	0.23	4.07	0.29	0.04	9.58	0.08

Table 3. Qualitative traits of Gladiolus hybrid selection IHRG-7 with check Pink Friendship

Sl. No.	Trait	IHRG -7	Pink Friendship
1.	Floret Type	Open-faced	Open-faced
2.	Floret texture	Medium	Medium
3.	Floret structure	Wavy	Wavy
4.	Floret placement	Good	Good
5.	Floret colour	Red-Purple (65.B) having Red-Purple (62.A) streaks with Red-Purple (67.B) splash	Red (50.D) having Red (51.C) margin and White (155.D) lines with Yellow (2.D) blotch

Data presented in Table 2 showed significant differences between hybrid selection IIHRG-7 and Pink Friendship (check) for corm and cormel characters. IIHRG-7 recorded significantly higher number of corms (1.46) than Pink Friendship (1.03); however, Pink Friendship recorded more number of cormels per corm (52.11) than IIHRG-7 (26.63). Significantly higher diameter of corm (6.75 cm), cormel (1.17 cm), weight of corm (72.44 g) and cormel (0.57 g) was recorded in IIHRG-7. Corm diameter and corm weight are important traits for producing quality spikes, with higher number of florets with bigger size. Sankari *et al.* (2012) and Safeena and Thangam (2019) reported that genotypes Thumbolina, Priscilla, Candyman, Arka Amar and Darshan were found superior for corm number, corm weight and corm diameter.

The qualitative traits of IIHRG-7 and Pink Friendship are given in Table 3. The IIHRG-7 has novel flower colour (RHS colour chart) as Red-Purple (65.B) having Red-Purple (62.A) streaks with Red-Purple (67.B) splash with variegated spikes.

On the perusal of the data presented in Table 4 significant differences are recorded between hybrid selection IIHRG-11 and Psittacinus hybrid (check) for most of the vegetative and floral characters, however, flowering duration and vase life were found non-significant. The hybrid selection IIHRG-11 recorded significantly early spike emergence (66.66 days) and flowering (76.65 days) in comparison to Psittacinus hybrid (check) (78.24 days and 89.245 days, respectively). Shaikat *et al.*, (2013) also reported early spike emergence in Applause and Peter Pears and early flowering in Priscilla and Peter Pears. Psittacinus hybrid recorded significantly maximum plant height (150.38 cm), spike length (120.43 cm) and rachis length (60.07 cm) than IIHRG-11 (120.72 cm, 95.18 cm and 48.81 cm, respectively). However, IIHRG-11 recorded significantly maximum floret diameter (9.46 cm), number of florets per spike (17.54) and florets remain open at a time (6.86) than Psittacinus hybrid (8.25 cm, 16.68 and 4.75, respectively), while, maximum total number of spikes per corm (3.92) and marketable spikes per corm (2.43) were recorded in Psittacinus hybrid than IIHRG-11 (1.92 and 1.70, respectively). The genotypes with more number of florets remain open at a time on the spike are more suited for exhibition purpose. The more number of spikes per corm are directly related to the higher productivity per unit area. Swaroop *et al.* (2018) evaluated 27 gladiolus

hybrids and reported that hybrids Suchitra x Melody and Green Pasture x Regency recorded maximum plant height, spike length and rachis length, while, hybrids Suchitra x Melody and Bindiya (mutant) recorded more number of florets per spike, whereas, hybrids Suchitra x Melody and Green Pasture x Regency recorded higher number of shoots per plant. However, Bhat *et al.* (2017) evaluated 60 genotypes of gladiolus for growth and flowering traits and recommended that genotypes Eurovision, Jester Gold, Priscilla, Vink's Glory, White Friendship etc. are best suited for cut flower under temperate conditions of Kashmir.

Data presented in Table 5 showed significant differences between hybrid selection IIHRG-11 and Psittacinus hybrid (check) for most of the corm and cormel traits except number of cormels per corm. Psittacinus hybrid recorded significantly higher number of corms per plant (3.64), diameter of cormel (1.78 cm) and weight of cormel (3.10 g) than IIHRG-11 (1.91, 1.53 cm and 1.08 g, respectively), whereas, IIHRG-11 recorded significantly higher corm diameter (6.64 cm) and corm weight (64.44 g) than Psittacinus hybrid (5.22 cm and 44.33 g, respectively). Bhat *et al.* (2017) evaluated 60 genotypes of gladiolus for corm and cormels traits and recommended genotypes Buff Beauty, Mayur, Priscilla, Pusa Suhagin, Regency etc. are best suited for corm production under temperate conditions of Kashmir.

The qualitative traits of IIHRG-11 and Psittacinus are given in Table 6. The IIHRG-11 has novel flower colour (RHS colour chart) as Red (41.C) having Red (41.A) margin. Blotch Red (46.B) with Yellow (13.C) border and have resistance to *Fusarium* wilt disease.

Data presented in Table 7 indicated that the hybrid selection IIHRG-11 recorded 18.52% disease incidence with zero per cent mortality which comes under resistant category, while, check Pink Friendship recorded 33.33% disease incidence with 18.52 per cent mortality which comes under moderately susceptible category as categorized by Shanmugam *et al.*, 2009.

On the basis of three years of evaluation, gladiolus hybrid selections IIHRG-7 was found to be promising for novel flower colour and variegated spike, and IIHRG-11 for novel flower colour and resistant to *Fusarium* wilt disease. These hybrid selections will be useful in developing new gladiolus hybrid selections with novel flower traits and resistant to *Fusarium* wilt disease.

Table 4. Vegetative and floral traits of Gladiolus hybrid selection IIHRG-11 with check Psittacinus hybrid (pooled data of three years)

Genotype	Days to spike emergence	Days to flower	Plant height (cm)	Spike length (cm)	Rachis length (cm)	Floret diameter (cm)	No. of florets per spike	Florets remain open at a time	No. of spikes/corn	No. of marketable spikes/corn	Flowering duration (days)	Vase life (days)
IIHRG-11	66.66	76.65	120.72	95.18	48.81	9.46	17.54	6.86	1.92	1.70	11.70	7.12
Psittacinus hybrid	78.24	89.24	150.38	120.43	60.07	8.25	16.68	4.75	3.92	2.43	11.96	7.00
C.D. at 5%	1.96	2.14	3.96	2.75	2.47	0.21	0.64	0.25	0.46	0.40	NS	NS

Table 5. Corm and cormel traits of Gladiolus hybrid selection IIHRG-11 with check Psittacinus hybrid (pooled data of three years)

Genotype	Corm per corm (Nos.)	Cormel per corm (Nos.)	Diameter of corm (cm)	Diameter of cormel (cm)	Weight of corm (g)	Weight of cormel (g)
IIHRG-11	1.91	10.14	6.64	1.53	64.44	1.08
Psittacinus hybrid	3.64	10.90	5.22	1.78	44.33	3.10
C.D. at 5%	0.28	NS	0.18	0.06	4.47	0.12

Table 6. Qualitative traits of Gladiolus hybrid selection IIHRG-11 with check Psittacinus hybrid

Sl. No.	Trait	IIHRG -11	Psittacinus hybrid
1.	Floret Type	Open-faced	Hooded
2.	Floret texture	Thick	Medium
3.	Floret structure	Slightly ruffled	Plain
4.	Floret placement	Double row	Fair
5.	Floret colour	Red (41.C) having Red (41.A) margin. Blotch Red (46.B) with Yellow (13.C) border	Red (39.A) with orange-Red (34.A) margin. Blotch Yellow (8.B)

Table 7. Disease incidence (%) and mortality (%) in IIHRG-11 with Pink Friendship (check) as influenced by *Fusarium* inoculum

Genotype	Disease incidence (%)	Mortality (%)
IIHRG-11	18.52 (18.47)	0.00 (2.87)
Pink Friendship	33.33 (30.95)	18.52 (18.47)
SEm±	3.46	3.14
CD (P=0.01)	20.63	18.70

Note: Values within parenthesis are *arc sign* transformed values

REFERENCES

- Anonymous. 2016. Horticulture crops estimates for the year 2015-16. National Horticulture Board, Haryana. Ministry of Agriculture and Farmer's Welfare, Government of India.
- Bhat, Z.A., Nazki, I.T., Nelofar and Hamid, B. 2017. Evaluation of gladiolus cultivars for growth, flowering, spike yield and corm yield under temperate regions of Kashmir. *Ind. Hort. J.*, **7**(3/4): 203-207.
- Elewa, I.S., Hilal, A.A., Hassan, S.E. and Abd El Malak, S. 2001. Studies on *Fusarium* disease (*Fusarium oxysporum* Schlecht. f. sp. gladioli (Massey) Synd. and Hans. of gladiolus. *Egyptian J. Phytopathol.*, **22**(2): 83-95.
- Kakade, D.S., Jadhav, S.B. and Katwate, S.M. 2016. Management of *Fusarium* wilt in Gladiolus. *Int. J. Pure App. Biosci.*, **4**(5): 127-132.
- Lakshman, D.K., Pandey, R., Kamo, K., Bauchan, G. and Mitra, A. 2012. Genetic transformation of *Fusarium oxysporum* f. sp. gladioli with *Agrobacterium* to study pathogenesis in Gladiolus. *Eur. J. Plant Pathol.*, **133**(3): 729-738.
- Massey, L.M. 1926. *Fusarium* rot of gladiolus corms. *Phytopathol.*, **52**: 567-572.
- Nelson P.E., Toussoun T.A. and Cook R.J. 1981. *Fusarium: Diseases, Biology, and Taxonomy*. Pennsylvania State University Press; University Park, Pennsylvania, USA.
- Panse, V.G. and Sukatme, P.V. 1967. Statistical methods for agricultural workers, Indian Council of Agricultural Research, New Delhi. p.155.
- Riaz, T., Khan, S.N. and Javaid, A. 2010. Management of corm-rot disease of gladiolus by plant extracts. *Nat. Prod. Res.*, **24**: 113-118.
- Safeena, S.A. and Thangam, M. 2019. Field performance of Gladiolus cultivars for growth, yield and quality cut flower production under humid agro-climatic conditions of Goa. *Int. J. Agr. Sci.*, **11**(3): 7797-7800.
- Sankari, A., Anand, M. and Arulmozhiyan, R. 2012. Evaluation of Gladiolus cultivars in Eastern Ghats of Tamil Nadu. *J. Hortl. Sci.*, **7**(2): 206-208.
- Shanmugam, V., Ajit, N.S., Ram, R., and Dhyani, D. 2009. Screening carnation and Gladiolus cultivars for vascular wilt resistance. *Indian Phytopathol.*, **62**(1): 117-118.
- Shaukat, S.A., Shah, S.Z.A., Shaukat, S.K. and Shoukat, S.W. 2013. Performance of gladiolus (*Gladiolus grandiflora* L.) cultivars under the climatic conditions of Bagh Azad Jammu and Kashmir Pakistan. *J. Cent. Eur. Agric.*, **14**(2): 158-167.
- Swaroop, K., Singh, K.P., Kumar, P. and S.S. Sindhu. 2018. Improvement and performance of gladiolus hybrids for flower traits/novel colour and higher corm multiplication. *Int. J. Agric. Innov. Res.*, **6**(4): 2319-1473.